DKB FARMS & SERVICES LLC 4945 MARATHON RD COLUMBIAVILLE, MI 48421 810-614-6099

ORG NIC MEDIUM RED CLOVER OT # WHIT-CLOV-12 NET WT. 50 LBS

CERT CRGANIC PROCESSOR # 4023H1011-10 Biodiversity Study Winter Wheat Red Clover Seeding Calculation

Three treatments BS(114, 217, 312, 405)
B8(118, 214, 310, 408) 1317(1/6,211,302,417)

Plot size 100'x30' = 3000 seft

· 3000 50# = 43,560 = . 67 Acres / 1 plot.

· 1 plot = . 07A @ . 07A x 12 (plots) = . 89 Acres.

Target Red Clover Seeding Rate 12-13 165/A • 12.5 165/A × . 84A = 10.5 165 sud.

4mph to feet / second.

- · 4mphx 5,280 ft = 21,126 ft/hr
- 21,120 ft/hr i 60 min = 352 ft/min 352 ft/min i 60 sec = 5.86 tt/sec

- 100' test strip i 5,86 fps = 17,06 sec

GANDY

Groms per min = Groms /Acre & mph x with (in)

- · width = 15' x 12" = 180" = 8 deflutors = 22-5"
- Grams pure Acre = 1215 x 454 grams = 544B grams A
- Groms or min = 5448 gp4 x 4 mph x 22.5" = 82.54 gpm = 42 2016 grams

 5940

 2016 grams

 pr 15544

	Torget 2	0.6 grams in	15 su	AVARO TO SECURITION OF THE SEC
	O	Bag		
Test		2	3	
l	16.6	16.74	16.64	
2	18,87	19.14	19.29	
3	21.6	21.1	21.23	
4	21.19	21.81	21.45	
5	20.74	20,81	20.65	
6	20,8	21.84	20,74	
		Aus		

= 20.796

≈ 2n a.

Added 2008 organ · 11.37 + 2.18 = 13.55 165.

Seeded Red Clover

· 13.55-3.17=10.38 165. 165 vsel.

March 18,2015

Area · 100 × 30 = 3000 ft =

· 3000 59 Ft : 435 60 59/14 = .0688 . . 0688 x 12 plots = 8000 Acres course.

· 10.38 lbs spread = . B26 Acres = 12.566 lbs / Acre

2015 Biodiversity Frost Seeding Wheat Plot Size 30' x 100' = 3000 sq ft 3000 \(\frac{4}{3}\)\(\frac{5}{60} = .068 3 treatments \(\chi \) 4 (eps = 12 plots 12 plots \(\chi \). 068 = .826 acres total Grams per acre = \frac{1216s}{A} \(\chi \) \(\frac{116s}{10s}\) Grams per min = \frac{5448apa \(\chi \) \(\frac{4mph \(\chi \) \(\chi \) \(\chi \)}{5940

82.54 ÷ 4 = 20,63 /15 sec grams per min grams in 15 sec

Tractor need to go 100ft in 17.06 sec

1. 17.96 2. 17.56 3. 16.83 4. 17.43

Grams per minute = Grans/Acre × MPH × width(in)
5946

2015 Protocol for Biodiversity Study LTER at Kellogg Biological Station, Michigan State University

System	Treatment	Plot Numbers			1	Crop Rotation
		Rep 1	Rep 2	Rep 3	Rep 4	
Α	B1	106	201	303	401	Fall Fallow
Α	B2	102	206	316	402	Spring Fallow
В	B 3	111	203	306	403	Corn covers A & C - Soybeans cover B - Wheat covers A & C
В	B 4	104	220	307	404	Soybeans over B - Wheat overs A & C - Corn overs* A & C
B	B5	114	217	312	405	Wheat covers A & c - Corn covers A & c - Soybeans cover B
С	B6	107	207	317	406	Corn over A - Soybeans - Wheat over A
С	B 7	105	219	305	407	Soybeans - Wheat over A - Corn over A
C	B8	118	214	310	408	Wheat cover A - Corn cover A - Soybeans
D	B9	119	205	314	409	Corn - Soybeans - Wheat
D	B10	117	209	320	410	Soybeans - Wheat - Corn
D	B11	110	216	309	411	Wheat - Corn - Soybeans
E	B12	109	202	313	412	Soybeans - Corn
E	B13	113	212	319	413	Corn – Soybeans
E	B14	115	204	304	414	Soybeans - Wheat
F	B15	112	213	301	415	Corn cover A - Corn cover A - Corn cover A
F	B16	101	210	308	416	Soybeans cover c - Soybeans cover c - Soybeans cover c
A FRANCE	B17	116	211	302	417	Wheat cover A - Wheat cover A - Wheat cover A
G	B18	108	208	311	418	Corn – Corn – Corn
G	B19	103	218	315	419	Soybeans - Soybeans - Soybeans
G	B20	120	215	318	420	Wheat - Wheat - Wheat
Н	B21	100	200	300	421	Continuous Fallow

*Cover A: Red Clover Cover B: Crimson Clover Cover C: Cereal Rye

System	Descriptions
Α	Fallow system: No crop is planted. Plots are tilled once a year.
В	One annual crop with two cover crops. Three year crop rotation.
C	One annual crop with one cover crop. Three year crop rotation.
D	One annual crop with no cover crop. Three year crop rotation.
E	One annual crop with no cover crop. Two year crop rotation.
F	One annual crop with one cover crop. Monoculture cropping system (no crop rotation).
G	One annual crop with no cover crop. Monoculture cropping system (no crop rotation).
H	Continuous fallow system: No cover, no crop growth. Plots are tilled as needed
	(2 - 6 times) a year to prevent plant growth from becoming established.

Research Objective: Incorporating biological diversity into weed management. Determine the impact of crop rotation and cover crops on weed communities in row crops.

Notes: All plots will be managed like the LTER main site treatment 4 plots.

No herbicides and no synthetic nitrogen will be used on any treatment.

This study was established in 2000. In 2000 and 2001 some treatments received fertilizer and herbicides.

Beginning in 2002 all treatments and plots have been treated like the LTER main site treatment 4, no herbicides and no synthetic fertilizer. Plot size = 30' x 90' (9 meters x 27meters).

This is a working protocol used for planning purposes. Due to potential changes in chemicals, fertilizer, varieties planted, planting dates etc...

please refer to the agronomic field log for actual field operations that take place during 2015.