







VARIETY: 25R40 LOT: H2GRA11009-LIT-1019 BATCH: 2477958 MATERIAL: 25R40-N341 TESTED: JULY 2015 NET WT: 50 LB (22.68 KG) KIND: WHEAT GERM: 90% ORIGIN: MI,OH

LIT1

PATENT PROTECTED

See Bag/Tag/Invoice for important terms of purchase and limited license. Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is prohibited.

Noxious Weed Seed Per Pound: None Other Crop Seed: 0.10% Pure Seed: 99.00% Inert Matter: 0.90% Weed Seed: 0.00%

10,500 Seeds / 14



CAUTION - This seed has been treated at manufacturers recommended rates with difenoconazole, mefenoxam, and sedaxane fungicides and imidacloprid insecticide. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice. Do not use for feed, food, or oil purposes. Slore away from feeds and toodstuffs. KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS. HAZARD COMMUNICATION DATA

CAUTION - TREATED SEED

Safety and Health: Corrosive. Causes irreversible eye damage. Wear long-sleeved shirt, long pants, shoes and socks, protective eyewear and chemical resistant gloves when handling treated seeds. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact skin or clothing, Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. For more information, call Pioneer at 1-800-342-7123. Restricted Entry Interval (REI): Do not enter or allow worker entry into treated areas during the REI of 48 hours. Exception: If the seed is treated with the predict and the treated area is a lifetime to the period. the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Environmental Hazards: This product is toxic to fish and highly toxic to birds and aquatic invertebrates. For terrestrial uses. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. If treated sed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect Intracted seeds spilled during loading. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Ground Water Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Restrictions: Green wheat or triticale forage may not be grazed or fed to livestock until 55 days after planting. Do not plant any crop other than those on the Dividend Extreme label within 30 days to fields in which treated seeds were planted. Vibrance: In the event of crop failure or harvest of a crop grown from Vibrance treated seed, the field may be replanted immediately to canola, soybean, cereal grains (barley, oat, rye, triticale, and wheat) corn, sorghum, crop group 6C (dried shelled pea and bean subgroup) and crop group 20A (rapeseed subgroup). All other crops may be planted no less than 30 days after planting treated seeds. Do not plant any crop other than corn, cereals, canola, soybeans, dry beans, dry peas, chickpeas and lentils, within 365 days to fields in which seed treated with Vibrance were planted. The maximum number of applications per season is 2. Imidacloprid restriction: Treated areas may be replanted with any crop specified on imidacloprid label, or any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. Areas planted with treated seed may be replanted immediately with artichoke, barley, Brassica (cole) leafy vegetables, borage, bult vegetables, canola, cilantro, corn (field), corn (pop), corn (sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, ground cherry, leaf petiole vegetables, leafy vegetables, legume vegetables (succulent or dried, including soybean), millet, mustard seed, oats, okra, pepinos, pepper, potato rapesed, rye, safflower, sorghum, soybean, strawberry, sugarbeet, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, wheat. These areas may also be replanted after 30 days with cereals, including buckwheat, rice. Do not plant any other crop in the treated area for at least one year after treated seeds are planted.

7/2014

Pioneer is a member of Excellence Through Stewardship[®] (ETS). Pioneer products are commercialized in accordance with ETS Product Launch Stewardship Guidance and in compliance with the Pioneer policies regarding stewardship of those products. Crops and materials containing biotech traits may only be exported to or used, processed, or sold in jurisdictions where all necessary regulatory approvals have been granted for those crops and materials. It is a violation of nationa and international laws to move materials containing biotech traits across borders into jurisdictions where their import is not permitted. Grovers should discuss these issues with their purchaser or grain handler to confirm the purchaser or handler's position on products being purchased. Excellence Through Stewardship⁶ is a registered trademark of the Biotechnology Industry Organization.

tered trademark of Pioneer Hi-Bred Internati © 2014 Pioneer Hi-Bred International, Inc. PIONEER HI-BRED INTERNATIONAL, INC P.O. Box 256 • Johnston, Iowa 50131-025 ®Registere tional, Inc









VARIETY: 25R40 LOT: H2GRA11014-GRA-1028 BATCH: 2506900 MATERIAL: 25R40-N341 TESTED: JULY 2015 NET WT: 50 LB (22.68 KG) KIND: WHEAT GERM: 90% ORIGIN: MI

GRA1

PATENT PROTECTED

See Bag/Tag/Invoice for important terms of purchase and limited license. Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is prohibited.

Noxious Weed Seed Per Pound: None Other Crop Seed: 0.10% Pure Seed: 99.00% Inert Matter: 0.90% Weed Seed: 0.00%

10,000 Seeds/15



CAUTION - This seed has been treated at manufacturers recommended rates with difenoconazole, metenoxam, and sedaxane fungicides and imidacloprid insecticide. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice. Do not use for feed, food, or oil purposes. Store away from feeds and toodstuffs. KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS. HAZARD COMMUNICATION DATA

CAUTION -- TREATED SEED

Safety and Health: Corrosive. Causes irreversible eye damage. Wear long-sleeved shirt, long pants, shoes and socks, protective eyewear and chemical resistant gloves when handling treated seeds. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact skin or clothing, Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. For more information, call Pioneer at 1-800-342-7123. Restricted Entry Interval (REI): Do not enter or allow worker entry into treated areas during the REI of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Environmental Hazards: This product is toxic to fish and highly toxic to birds and aquatic invertebrates. For terrestrial uses. Do not apply directly to water, to a reas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Ground Water Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Restrictions: Green wheat or triticale forage may not be grazed or fed to livestock until 55 days after planting. Do not plant any crop other than those on the Dividend Extreme label within 30 days to fields in which treated seeds were planted. Vibrance: In the event of crop failure or harvest of a crop grown from Vibrance treated seed, the field may be replanted immediately to canola, soybean, cereal grains (barley, oat, rye, triticale, and wheat) corn, sorghum, crop group 60 (dried shelled pea and bean subgroup) and crop group 20A (rapeseed subgroup). All other crops may be planted no less than 30 days after planting treated seeds. Do not plant any crop other than corn, cereals, canola, soybeans, dry beans, dry peas, chickpeas and lentils, within 365 days to fields in which seed treated with Vibrance were planted. The maximum number of applications per season is 2. Imidacloprid restriction: Treated areas may be replanted with any crop specified on imidacloprid label, or any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. Areas planted with treated seed may be replanted immediately with artichoke, barley, Brassica (cole) leafy vegetables, borage, bulb vegetables, canola, cilantro, corn (field), corn (pop), corn (sweet), cotton, cranberry, crambe, vegetables, carlos, charlos, com (reac), com (pop), com (erec), cocho, carlos (reach), coch least one year after treated seeds are planted.

7/2014

Pioneer is a member of Excellence Through Stewardship[®] (ETS). Pioneer products are commercialized in accordance with ETS Product Launch Stewardship Guidance and in compliance with the Pioneer policies regarding stewardship of those products. Crops and materials containing biotech traits may only be exported to or used, processed, or sold in jurisdictions where al necessary regulatory approvals have been granted for those crops and materials. It is a violation of nationa Indicatory registrational laws to move materials containing biotech trails across borders into jurisdictions where their import is not permitted. Growers should discuss these issues with their purchaser or grain handler to confirm the purchaser or handler's position on products being purchased. Excellence Through Stewardship' is a registered trademark of the Biotechnology Industry Organization.

®Registered trademark of Pioneer Hi-Bred International, Inc. © 2014 Pioneer Hi-Bred International, Inc. PIONEER HI-BRED INTERNATIONAL, INC. P.O. Box 256 - Johnston, Iowa 50131-0256



52840-N241 5204044



VARIETY: 25R40 LOT: G2MCL11002-00-0083 BATCH: 2594044 MATERIAL: 25R40-N541 TESTED: SEPT 2015 NET WT: 50 LB (22.68 KG)

KIND: WHEAT GERM: 80% ORIGIN: IL

MCL1

PATENT PROTECTED

See Bag/Tag/Invoice for important terms of purchase and limited license Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is prohibited.

Noxious Weed Seed Per Pound: None Other Crop Seed: 0.10% Pure Seed: 99.00% Inert Matter: 0.90% Weed Seed: 0.00%

11,000 Seeds/16



CAUTION - This seed has been treated at manufacturers recommended rates with difenoconazole, metenoxam, thiabendazole, and sedaxane fungicides and imidacloprid insecticide. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs. KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS. HAZARD COMMUNICATION DATA

CAUTION -- TREATED SEED

Salety and Health: Corrosive. Causes irreversible eye damage. Wear long-sleeved shirt, long pants, shoes and socks, protective eyewear and chemical resistant gloves when handling treated seeds. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. For more information, call Pioneer at 1-800-342-7123. Restricted Entry Interval (REI): Do not enter or allow worker entry into treated areas during the REI of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Environmental Hazards: This product is toxic to fish and highly toxic to birds and aquatic invertebrates. For terrestrial uses. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Ground Water Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Restrictions: Green wheat or triticale forage may not be grazed or fed to livestock until 55 days after planting. Do not plant any crop other than those on the Dividend Extreme label within 30 days to fields in which treated seeds were planted. Vibrance: In the event of crop failure or harvest of a crop grown from Vibrance treated seed, the field may be replanted immediately to canola, soybean, cereal grains (barley, oat, rye, triticale, and wheat) corn, sorghum, crop group 6C (dried shelled pea and bean subgroup) and crop group 20A (rapeseed subgroup). All other crops may be planted no less than 30 days after planting treated seeds. Do not plant any crop other than corn, cereals, canola, soybeans, dry beans, dry peas, chickpeas and lentils, within 365 days to fields in which seed treated with Vibrance were planted. The maximum number of applications per season is 2. Imidacloprid restriction: Treated areas may be replanted with any crop specified on imidacloprid label, or any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. Areas planted with treated seed may be replanted immediately with artichoke, barley, Brassica (cole) leafy vegetables, borage, bulb vegetables, canola, cilantro, corn (field), corn (pop), corn (sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, ground cherry, leaf petiole vegetables, leafy vegetables, legume vegetables (succulent or dried, including soybean), millet, mustard seed, oats, okra, pepinos, pepper, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeet, sunflower, tomatilib, tomato, triticale, root and tuber vegetables, watercress, wheat. These areas may also be replanted after 30 days with cereals, including buckwheat, rice. Do not plant any other crop in the treated area for at least one year after treated seeds are planted.

7/2014

1

Pioneer is a member of Excellence Through Stewardship® (ETS). Pioneer products are commercialized in accordance with ETS Product Launch Stewardship Guidance and in compliance with the Pioneer policies regarding stewardship of those products. Crops and materials comparate with the Photeer poinces regarding stewardship of those products. Crops and materials containing blotch traits may only be exported to or used, processed, or sold in jurisdictions where all necessary regulatory approvals have been granted for those crops and materials. It is a violation of national and international laws to move materials containing biotech traits across borders into jurisdictions where their import is not permitted. Growers should discuss there is sues with their purchaser or grain handler to confirm the purchaser or handler's position of products being purchased. Excellence Through Stewardship® is a registered trademark of the Biotechnology Industry Organization.

M.M.Trademarks and service marks of DuPont, Pioneer or their respective owners © 2015 PHII.
© 2015 Pioneer Hi-Bred International, Inc.
PIONEER HI-SRED INTERNATIONAL, INC., PO Box 256, Johnston, IA 50131-0256



VARIETY: 25R40 LOT: G2GRA11008-00-0165 BATCH: 2593630 MATERIAL: 25R40-N541 TESTED: SEPT 2015 NET WT: 50 LB (22.68 KG) KIND: WHEAT GERM: 80% ORIGIN: MI

GRA1

PATENT PROTECTED

See Bag/Tag/Invoice for important terms of purchase and limited license. Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is prohibited.

Noxious Weed Seed Per Pound: None Other Crop Seed: 0.10% Pure Seed: 99.00% Inert Matter: 0.90% Weed Seed: 0.00% 10,000 Seeds IL IONEE

CAUTION - This seed has been treated at manufacturers recommended rates with difenoconazole, metenoxam, thiabendazole, and sedaxane fungicides and imidacloprid insecticide. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice. Do not use for feed, food, or oil purposes. Store away from feeds and toodstutts, KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS HAZARD COMMUNICATION DATA

CAUTION - TREATED SEED

Safety and Health: Corrosive. Causes irreversible eye damage. Wear long-sleeved shirt, long pants, shoes and socks, protective eyewear and chemical resistant gloves when handling treated seeds. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. For more information, call Pioneer at 1-800-342-7123. Restricted Entry Interval (REI): Do not enter or allow worker entry into treated areas during the REI of 48 hours. Exception: If the sed is treated with the product and the treated sed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Environmental Hazards: This product is toxic to fish and highly toxic to birds and aquatic invertebrates. For terrestrial uses. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading. Dispose of all excess treated seed. Dispose of seed packaging in accordance with local requirements. Ground Water Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Restrictions: Green wheat or triticale forage may not be grazed fed to livestock until 55 days after planting. Do not plant any crop other than those on the or Dividend Extreme label within 30 days to fields in which treated seeds were planted. Vibrance: In the event of crop failure or harvest of a crop grown from Vibrance treated seed, the field may be replanted immediately to canola, soybean, cereal grains (barley, oat, rye, triticale, and wheat) corn, sorghum, crop group 6C (dried shelled pea and bean subgroup) and crop group 20A (rapeseed subgroup). All other crops may be planted no less than 30 days after planting treated seeds. Do not plant any crop other than corn, cereals, canola, soybeans, dry beans, dry beas, chickpeas and lentils, within 365 days to fields in which seed treated with Vibrance were planted. The maximum number of applications per season is 2. Imidacloprid restriction: Treated areas may be replanted with any crop specified on imidacloprid label, or any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. Areas planted with treated seed may be replanted immediately with artichoke, barley. Brassica (cole) leafy vegetables, borage, bulb vegetables, canola, cilantro, corn (field), corn (pop), corn (sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, ground cherry, leaf petiole vegetables, leafy vegetables, legume vegetables (succulent or dried, including soybean), millet, mustard seed, oats, okra, pepinos, pepper, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeet, sunflower, tornatilio, tomato, triticale, root and tuber vegetables, watercress, wheat. These areas may also be replanted after 30 days with cereals, including buckwheat, rice. Do not plant any other crop in the treated area for at least one year after treated seeds are planted.

7/2014

Pioneer is a member of Excellence Through Stewardship® (ETS).

Pioneer is a member of Excellence Through Stewardship[®] (ETS). Pioneer products are commercialized in accordance with ETS Product Launch Stewardship Guidance and in compliance with the Pioneer policies regarding stewardship of those products. Crops and materials containing biotech traits may only be exported to or used, processed, or sold in jurisdictions where all necessary regulatory approvals have been granted for those crops and materials. It is a violation of national and international laws to move materials containing biotech traits across borders into jurisdictions where their import is not permitted, Growers should discuss these issues with their purchaser or grain handler to confirm the purchaser or thandler's position on products being purchased. Excellence Through Stewardship[®] is a registered trademark of the Biotechnology Industry Organization.

SM Intrademarks and service marks of DuPont, Pioneer or their respective owners @ 2015 PHII. @ 2015 Primeer HI-Bred Imemational, Inc. PIONEER HI-BRED INTERNATIONAL, INC., PO Box 256, Johnston, IA 50131-0256