

12195771 -- 80.000 SEEDS/UNIT-- A500 -- AR
(01) 0 0883580952682 (10) 757XMM3JM 1757



Hybrid Corn Seed Blend

DKC52-84RIB

DEKALB BRAND
(GENSSRIB)

RELATIVE MATURITY: OVERALL -102

GROWING DEGREE UNITS MID-POLLINATION: 1266

BLACK LAYER: 2550

LOT NO. 757XMM3JM

ORIGIN GERM DATE TESTED

VARIETY 1052505: 94.00% NE 95% 11/15

TREATMENT: WAIH2V

VARIETY 1056251 5.00% MI 95% 11/15

TREATMENT: WAIH2V

INERT MATTER: 0.40%

WEED SEED: 0.00%

OTHER CROP SEED: 0.60%

NOXIOUS WEEDS/LB: NONE

KIND: FIELD CORN

SUGGESTED PLATE:

JD B25

CIH C25



This is a product of MONSANTO's reasearch program offering unique genetic characteristics forspecific grower needs and protected by U.S. patent(s) : PENDING

genuity

SMARTSTAX[®] CORN

RIB COMPLETE[®]

SEE INSIDE FOR IRM INFORMATION

A500

OPEN
HERE

CODE SEED-APPLIED TECHNOLOGY

A	Metalaxyl
D	Azoxystrobin
H1	Clothianidin 250
H2	Clothianidin 500
H3	Clothianidin 1250
I	Trifloxystrobin
M	Fludioxonil
P	Penflufen
T	Torque®
V	<i>Bacillus firmus I-1582</i>
W	Iproconazole
X	Mefenoxam



INSECT DISEASE CORN

Seed Treatment Products

WARNING. TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS.

This seed has been treated at the manufacturer's recommended rate with the products identified by code on the front of this seed tag, and are decoded in the table.

General Information: Handle treated seeds in ventilated areas. Avoid breathing or making contact with dust. If contaminated, remove clothing and launder before reuse. Wash hands and face before eating, drinking, or smoking. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect spilled treated seeds spilled during loading and planting.

Personal Protective Equipment: Wear long sleeved shirt, long pants, socks and shoes, chemical resistant gloves, and, if indicated for the specific product, goggles and a dust mask when handling treated seeds.

Hazards to Humans and Animals: Harmful to humans if swallowed, inhaled or absorbed through skin. Ingesting of treated seed may be hazardous to domestic animals, livestock and other wildlife.

Disposal: Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements. Do not use empty seed bags for any other purpose. Do not contaminate water bodies when disposing of planting equipment washwaters. Dispose of washwaters, seed packaging or containers in accordance with local requirements. Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.

FIRST AID: Eyes: Hold eye open and rinse gently with water for 15 minutes. If wearing contact lenses, remove after first 5 minutes and continue rinsing. **Skin or clothing:** Remove contaminated clothing. Rinse skin immediately with plenty of water. Use soap if available. **Swallowed:** Call poison center or doctor immediately for treatment advice. Sip water if able to swallow. Do not induce vomiting unless told to do so by the poison center or doctor. Do not give anything by mouth to an unconscious person. **Inhaled:** Move to fresh air. If not breathing, call 911 or ambulance, and then give artificial respiration. Call poison center or doctor for further treatment advice. **FOR EMERGENCY MEDICAL TREATMENT INFORMATION, CALL COLLECT DAY OR NIGHT: (314) 694-4000.**

PRODUCT SPECIFIC ADDITIONAL INFORMATION. CLOTHIANIDIN-TREATED SEED: Immediate Plant-Back: Cereal grains, field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetable. **30-Day Plant-Back:** Grasses, non-grass animal feeds, dry beans; **8-Month Plant-Back:** Sugar cane; **12-Month Plant-Back:** Any crops without an earlier plant-back interval.

CLOTHIANIDIN + BACILLUS FIRMUS I-1582-TREATED SEED: This seed has been treated with PONCHO®/VOTIVO®, which contains 40.3% clothianidin and 8.1% *Bacillus firmus I-1582*. Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch. Regardless of type of application (seed treatment or foliar), do not apply more than 0.2 lbs (Soybean), 0.1 lbs (corn), 0.2 lbs (cotton), 0.14 lbs (sugarbeet), or 0.02 lbs (sorghum) of active ingredient clothianidin per acre per season. This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds. This compound is toxic to birds and mammals. **Immediate Plant-back:** Cereal grains, field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetable; **30-Day Plant-back:** Grasses, non-grass animal feeds, dry beans; **8-Month Plant-back:** Sugar cane; **12-Month Plant-back:** Any crops without an earlier plant-back interval.

FLUDIOXONIL + MEFENOXAM-TREATED SEED: Corn and sorghum forage may not be grazed until 30 days after planting.

PENFLUFEN-TREATED SEED: After the seeds and/or seed pieces have been planted in soil or other planting media, DO NOT enter or allow worker entry into areas with treated seeds during the restricted-entry interval of 12 hours, except that workers may enter the areas with treated seeds without restriction if there will be no worker contact with the soil/media subsurface. Treated seeds must be planted at a minimum depth of 0.5 inches. Crops for which there are registered uses of penflufen, which includes alfalfa, potatoes (Crop Subgroup 1C), legume vegetables (Crop Group 6/7), cereal grains (Crop Group 15/16), and oil seeds (Crop Group 20) may be replanted at any time. All other crops may be planted no less than 30 days after planting treated seeds. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. The maximum number of application per year is 3. For corn: Do not exceed 0.0033 lb ai/A (1.5 g ai/A) per crop season.

TRIFLOXYSTROBIN-TREATED SEED: After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours. **Exception:** Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media. Areas planted with treated seed may be replanted immediately with any crop for which a tolerance exists for trifloxystrobin. Do not plant any crop without trifloxystrobin tolerances until 30 days after planting trifloxystrobin-treated seed.

Acceleron and Design® is a registered trademark of Monsanto Technology LLC. Poncho® and VOTIVO® are registered trademarks of Bayer. Torque® is a registered trademark of Novozymes A/S.

Call 1-800-8 DEKALB (1-800-833-5252) with any questions, feedback or additional seed needs.

NOTICE: See bag for limitation of warranties, liability and remedies.

Monsanto Company - Attention Seed Box - 800 N. Lindbergh Blvd. - St. Louis, MO 63167

NET WEIGHT PRINTED ON BAG

40088M7-61/53

©2008-2014 MONSANTO COMPANY



40088M7-61/53