

CAUTION FEED, FOOD, ETHANOL OR OIL PURPOSES, STORE Refill bag or container with feed or foodst HAZARD COMMUNICATION DATA TREATED SEED CORN - CAUTION

PRODUCT USE STATEMENT the manufacture is (Bt) Crynn, he YIELDGARD® con-ctection or suppression lack cutworm, fall an lack cutworm, fall an lack cutworm, fall an larval

location, pest population, environmental conditions, and agricultural practices. These proteins and the genetic material necessary for their production in corn are registered under EPA Reg. No. 29964-11 YOU MUST SIGN A TECHNOLOGY AGREEMENT, READ THE PRODUCT USE GUIDE PRIOR TO PLANTING AND FOLLOW INSECT RESISTANCE MANAGEMENT (IRM) REQUIREMENTS. PATENT STATEMENT: The Herculex I and Herculex RW Insect Resistance technologies incorporated into these seeds are pre-acted under one or more U.S. patents. The purchase of these seeds includes a limited licence to production or development of a hybrid or different variety of seed is strictly proliabiled *YIELDGARD is a registered trademark used under license from Monsanto Co. Herculex® Insect Protection technology by Daw AgroSciences and Pioneer Hi-Bred, *Herculex is a registered trademark of Dow AgroSciences LLC. US AMX Rev 6-15.

AL 329 LICATIONS OF INCOMPATIBLE HERBICIDES

ACQUIT 15 UNDER AN AGREEMENT

Pioneer® brand products and conditions of purchase Pont, Pioneer DuPont, Pioneer 5 PHIL 6, Johnston, IA 50131-0256

12-2692-A OAMXtra 30966



SOUTHERN CORN/COTTON GROWING AREAS MUST FOLLOW THE REFUGE REQUIREMENTS DESCRIBED BELOW AND IN THE PRODUCT USE GUIDE PRIOR TO PLANTING. NO ADDITIONAL REFUGE IS NEEDED IN THE NON-COTTON GROWING AREAS.

- Refuge must be at least 20% and must be planted within 1/2 mile of the Bt corn borer resistant hybrid
- Refuge block or strips must be ≥ 4 rows
- Refuge hybrid must not express a Bt protein active against corn borer
- NON-COTTON GROWING AREAS: Foliar insecticide treatments for control of European corn borer, corn earworm, southwestern corn borer, fall armyworm, black cutworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, and sugarcane borer may be applied only if economic thresholds are reached for one or more of these target pests. Foliar insecticide treatments are also permitted for control of corn rootworm adults if economic thresholds are reached.

SOUTHERN CORN/COTTON GROWING AREAS: Foliar insecticide treatments for control of European corn borer, corn earworm, southwestern corn borer, fall armyworm, black cutworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, and sugarcane borer may be applied only if economic thresholds are reached for one or more of these target pests. In addition, the refuge can be protected from CRW damage by an appropriate seed treatment or soil insecticide; however, insecticides labeled for adult CRW control must be avoided in the refuge during the period of CRW adult emergence.

Microbial Bt insecticides must not be applied to non-Bt corn refuge plants.



Optimum® AcreMax® Xtra Insect Protection REFUGE OPTIONS



Cotton-Growing Areas are all the counties in: AR, LA, MS, AL, GA, FL, SC, NC and in:

MO counties of Dunklin, New Madrid, Pemiscot, Scott and Stoddard

OK counties of Beckham, Caddo, Comanche, Custer, Greer, Harmon, Jackson, Kay, Kiowa, Tillman and Washita

TN counties of Carroll, Chester, Crockett, Dyer, Fayette, Franklin, Gibson, Hardeman, Hardin, Haywood, Lake, Lauderdale, Lincoln, Madison, Obion, Rutherford, Shelby and Tipton

TX counties except Carson, Dallam, Hansford, Hartley, Hutchinson, Lipscomb, Moore, Ochiltree, Roberts and Sherman

VA counties of Dinwiddie, Franklin City, Greensville, Isle of Wight, Northampton, Southampton, Suffolk City, Surrey and Sussex