



DKC52-84RIB DEKALB BRAND (SSRIB)

RELATIVE MATURITY: OVERALL -102 GROWING DEGREE UNITS MID-POLLINATION: 1266

BLACK LAYER: 2550

LOT NO. 748YKH9JX

ORIGIN GERM DATE TESTED VAR ETY 1052505: 94.00% MI TREATMENT: FALH2V VARIETY 1056251 5.00 % MI TREATMENT: FALH2V INERT MATTER: 0. 0.40 % WEED SEED: 0.00 % OTHER CROP SEED: 0.60 % NOXIOUS WEEDS/LB:NONE KIND: FIELD CORN SUGGESTED PLATE:



JD B9 CIH C9

This is a product of MONSANTO's research program offering unique genetic characteristics for specific grower needs and protected by U.S. patent(s) : PENDING




40088P13-61 SEED-APPLIED TECHNOLOGY Metalaxyl CELERON Acceleron® B-300 SAT

F Prothioconazole H1 Clothianidin 250 H2 Clothianidin 500 Clothianidin 1250 H3 Trifloxystrobin Fluoxastrobin Bacillus firmus I-1582 V **Ipconazole** W

CODE

A

B



SEE INSIDE PANEL FOR TREATMENT SPECIFIC ADDITIONAL INFORMATION.

General Information: Handle treated seeds in ventilated areas. Avoid breathing or making contact with dust. If contaminated, remove clothing and launder before reuse. Wash hands and face before eating, drinking, or smoking.

Personal Protective Equipment: When opening this bag or loading, pouring, or handling the treated seed or seed pieces, wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material.

Hazards to Humans and Animals: Harmful to humans if swallowed, inhaled or absorbed through skin. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. After the seeds and/or seed pieces have been planted in soil or other planting media, DO NOT enter or allow worker entry into areas with treated seeds during the restricted-entry interval of 12 hours, except that workers may enter the areas with treated seeds without restriction if there will be no worker contact with the soil/media subsurface.

Storage and Disposal: Store treated seed away from food and feed, and do not allow access by children, pets, or livestock. Do not reuse or refill empty seed bags. Surplus treated seed or empty treated seed containers should be stored or disposed according to local requirements. Dispose of all excess treated seed. Do not contaminate bodies of water when disposing of excess treated seed or wash waters of planting equipment. Dispose of them in accordance with local requirements. Excess treated seed may be used for ethanol production only if: (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticides remain in ethanol by-products that are used in agronomic practice.

FIRST AID: Eyes: Hold eye open and rinse gently with water for 15 minutes. If wearing contact lenses, remove after first 5 minutes and continue rinsing. Skin or clothing: Remove contaminated clothing. Rinse skin immediately with plenty of water. Use scap if available. Swallowed: Call poison center or doctor immediately for treatment advice. Sip water if able to swallow. Do not induce vomiting unless told to do so by the poison center or doctor. Do not give anything by mouth to an unconscious person. Inhaled: Move to fresh air. If not breathing, call 911 or ambulance, and then give artificial respiration. Call poison enter or doctor for further treatment advice. FOR EMERGENCY MEDICAL TREATMENT INFORMATION, CALL COLLECT DAY OR NIGHT: (314) 694-4000.

PRECAUTIONARY INFORMATION CONTINUED ON INSIDE PANEL.

Call 1-800-8 DEKALB (1-800-833-5252) with any questions, feedback or additional seed needs. NOTICE: See bag for limitation of warranties, liability and remedies.

Monsanto Company - Attention Seed Box - 800 N. Lindbergh Blvd. - St. Louis, MO 63167 **NET WEIGHT PRINTED ON BAG**

40088P13-61

©2008-2016 MONSANTO COMPANY

