

### ABOVE/BELOW



## PIONEER.

BLEND/MIXTURE COMPARATIVE RM 91



P9188AMX-NM07 (10)700638

H3CON11052-0 OCT 2016 Field F14 NO: KIND: LOT Corn

SIZE: F14 ALG1 KERNELS/UNIT:80,000 FERTILE CYTOPLASM TESTED:

Hybrid:P9188YXR(90% of Pure Seed) Germ:95% Origin:MI

Hybrid: IRC95-01LR(10% of Pure Seed) Germ: 95% Origin:MI

The seed is a Mixture under the state laws of AL, AK, AZ, CT, DE, FL, GA, ID, IL, KS, LA, KY, MA, MD, MS, NC, NE, NJ, NV, NY, OK, OR, TN, UT, WV and WI.

See Bag/Tag/Invoice for important terms of purchase and limited license. Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is

Noxious Weed Per Pound: None Other Crop Seed: None Pure Seed: 99.5% Inert Matter: 0.5% Weed Seed: None















NOTICE: See important Product Use Information on back of tag.

CAUTION: This seed is treated at the manufacturer's recommended rates with Fludioxonil, Mefenoxam, Thiabendazole and Azoxystrobin fungicides, Thiamethoxam and Clothianidin insecticides and Bocillus Firmus. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT USE TREATED SEED FOR FEED, FOOD, ETHANOL OR OIL PURPOSES. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT REFILL BAG OR CONTAINER WITH FEED OR FOODSTUFF.

FEED, FOOD, ETHANOL OR OIL PURPOSES. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT REFILL BAS OR CONTAINER WITH FEED OR FOODSTUFF.

HAZARD COMMUNICATION DATA

TREATED SEED CORN — CAUTION

Salety & Health: Wear long-sleeved shirt, long pants, eye protection, shoes, socks and waterproof gloves when handling treated seeds. Treated seed and the dust from treated seed may be harmful if swallowed, inhaled or absorbed through the skin so take the proper precautions to prevent contact. Wash clothing separately from other dothers, in case of emergency, contact provider at 800-342-7128. Environmental Hazards. Dispose of all wocoss treated excell. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. This compound may be highly toxic to here seposed directly contact. Ensure that planting requirements in Intendening proceed or planting equipment wash waters. Dispose of seed packaging in accordance with local requirements. This compound may be highly toxic to here seposed directly contact. Ensure that planting requirements in Intendening proceed with manufacturents personal commendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds. This compound is toxic to birds and manufacts. Treated seed exposed of not oil surface may be hazardous to birds and manumads. Cover or collect treated seeds spilled during loading. Restrictions may be hazardous to birds and manumads. Cover or collect treated seed spilled during loading. Restrictions on brage may not be grazed until 30 days after planting. Treated seed segment on tool surface may be hazardous to graze and the spilled during loading. Restrictions on brage may not be grazed until 30 days after planting. Treated seed one for loading planting this until 20 to 10 spilled planting to the spilled during loading. Restrictions on brage may not be grazed until 30 days after planting. Treated seed one for loading planting tool to planting this manufact

PRODUCT USE STATEMENT: This seed is a blend of 10% refuge seed and 90% seed containing the Herculex® XTRA Insect Protection genes that produce a Bacillus thuringiensis (Bt) Cry1F protein and the Bt Cry34Ab1 and Cry35Ab1 proteins and also contains the YIELDGARD® Corn Borer gene that produces a Bt Cry1Ab protein that provide protection or suppression against susceptible European corn borer, southwestern corn borer, black cutworm, fall armyworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, sugarcane borer, stalk borer, and corn earworm; and also provide protection from larval injury caused by susceptible western corn rootworm, northern corn rootworm, and Mexican corn rootworm. Product responses may vary by location, pest population, environmental conditions, and agricultural practices. These proteins and the genetic material necessary for their production in corn are registered under EPA Reg. No. 29964-11
YOU MUST SIGN A TECHNOLOGY AGREEMENT, READ THE PRODUCT USE GUIDE PRIOR TO PLANTING AND FOLLOW INSECT RESISTANCE MANAGEMENT (IRM) REQUIREMENTS.

PATENT STATEMENT: The Herculex I and Herculex RW Insect Resistance technologies incorporated into these seeds are protected under one or more U.S. patents. The purchase

PATEMT STATEMENT: The Herculex I and Herculex RW Insect Resistance technologies incorporated into these seeds are protected under one or more U.S. patents. The purchase of these seeds includes a limited license to produce a single corn crop in the United States (or other applicable country). The use of seed from such a crop or the progeny thereof for propagation or seed multiplication or for production or development of a hybrid or different variety of seed is strictly prohibited.

"YIELDGARD is a registered trademark used under license from Monsanto Co. Herculex® Insect Protection technology by Dow AgroSciences and Pioneer HI-Bred.

"Herculex is a registered trademark of Dow AgroSciences LLC.

US AMX Rev. 6-15

PRODUCT USE STATEMENT: This hybrid contains the Roundup Ready® Corn 2 gene developed by Monsanto Company. The Roundup Ready gene provides resistance to glyphosate, the active ingredient in labeled Roundup® branded herbicides. WARNING: The Roundup Ready gene will safeguard this hybrid OnLY against applications of glyphosate, the active ingredient in labeled Roundup branded herbicides, when applied at load rates. The Roundup Ready gene WILL NOT safeguard this hybrid against applications of other herbicides which require a different herbicide resistance gene. Always read and follow herbicide label directions prior to use. ACCIDENTAL APPLICATIONS OF INCOMPATIBLE HERBICIDES TO THIS HYBRID COULD RESULT IN TOTAL REPLICATIONS. ates. The the equire a different ACCIDENTAL APP

PRODUCT USE STATEMENT: This hybrid contains the LibertyLink® gene developed by Bayer. These seeds and the plants grown from these seeds produce a PAT (phosphinothricin acatyltransferase) protein that provides resistance to the Liberty Herbicide (glufosinate). WARNING: The LibertyLink gene will safeguard this report of 0013 against applications of LibertyLink gene will safeguard this hybrid of 0013 against applications of LibertyLink gene will safeguard this hybrid directions prior to use. ACIDIENTAL APPLICATIONS OF INCOMPATIBLE HERBICIDES TO THIS HYBRID COULD RESULT IN TOTAL CROP LOSS.

HESULI IN TOTAL CROP LOSS.

The herbicide resistance technology contained in this seed is protected at least under one or more of following U.S. patent(s): RE44,962. The conditional purchase of these seeds conveys no license under patents to use these seeds, except to produce a single commercial crop (one time use only) in the UT States, and harvesting the grain solely for purposes of tood or feed applications or for industrial process in no event shall this seed be used in research or in breeding or for production of planting seed. LibertyLink® and the Water Droplet Design are trademarks of Bayer.

LL Corn Rev. 4-16



12-2692-A OAMXtra 30966

SOUTHERN CORN/COTTON GROWING AREAS MUST FOLLOW THE REFUGE REQUIREMENTS DESCRIBED BELOW AND IN THE PRODUCT USE GUIDE PRIOR TO PLANTING. NO ADDITIONAL REFUGE IS NEEDED IN THE NON-COTTON GROWING AREAS.

- Refuge must be at least 20% and must be planted within 1/2 mile of the Bt corn borer resistant hybrid
- Refuge block or strips must be ≥ 4 rows
- Refuge hybrid must not express a Bt protein active against corn borer
- NON-COTTON GROWING AREAS: Foliar insecticide treatments for control of European corn borer, corn earworm, southwestern corn borer, fall armyworm, black cutworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, and sugarcane borer may be applied only if economic thresholds are reached for one or more of these target pests. Foliar insecticide treatments are also permitted for control of corn rootworm adults if economic thresholds are reached.

SOUTHERN CORN/COTTON GROWING AREAS: Foliar insecticide treatments for control of European corn borer, corn earworm, southwestern corn borer, fall armyworm, black cutworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, and sugarcane borer may be applied only if economic thresholds are reached for one or more of these target pests. In addition, the refuge can be protected from CRW damage by an appropriate seed treatment or soil insecticide; however, insecticides labeled for adult CRW control must be avoided in the refuge during the period of CRW adult emergence.

Microbial Bt insecticides must not be applied to non-Bt corn refuge plants.

# In-field Refuge Separate Field Refuge Block Strips Optimum® AcreMax® Xtra Insect Protection ≥ 20% Corn Borer Refuge (e.g. non-Bt, herbicide tolerant, HXRW)

For additional refuge planning tools, you may also log onto **www.irmcalculator.com**.



20 % REFUGE Cotton-growing Areas



Cotton-Growing Areas are all the counties in: AR, LA, MS, AL, GA, FL, SC, NC and in:

MO counties of Dunklin, New Madrid, Pemiscot, Scott and Stoddard

OK counties of Beckham, Caddo, Comanche, Custer, Greer, Harmon, Jackson, Kay, Kiowa, Tillman and Washita

TN counties of Carroll, Chester, Crockett, Dyer, Fayette, Franklin, Gibson, Hardeman, Hardin, Haywood, Lake, Lauderdale, Lincoln, Madison, Obion, Rutherford, Shelby and Tipton

TX counties except Carson, Dallam, Hansford, Hartley, Hutchinson, Lipscomb, Moore, Ochiltree, Roberts and Sherman

VA counties of Dinwiddie, Franklin City, Greensville, Isle of Wight, Northampton, Southampton, Suffolk City, Surrey and Sussex