

12588263 -- 80,000 SEEDS/UNIT--  
-- AF (01) 0 0888346315725 (10) 748MMF9JX 1748



# DKC51-38RIB

DEKALB BRAND  
(SSRIB)

RELATIVE MATURITY: OVERALL -101

GROWING DEGREE UNITS MID-POLLINATION: 1275			
BLACK LAYER: 2525			
LOT NO. 748MMF9JX		ORIGIN GERM DATE TESTED	
VARIETY 1063249:	94.00% MI	95%	11/17
TREATMENT: FALH2VB			
VARIETY 1048633	5.00% MI	95%	11/17
TREATMENT: WAIH2V			
INERT MATTER:	0.40%		
WEED SEED:	0.00%		
OTHER CROP SEED:	0.60%		
NOXIOUS WEEDS/LB:	NONE		
KIND: FIELD CORN			
SUGGESTED PLATE:			
JD B9			
CIH C9			
Acceleron B-300 SAT Batch#:1735JS2-10Not Claimed			
Effective After:11/01/2019 For storage Requirements			
see Product-Specific Additional information on this label.			
This is a product of MONSANTO's research program offering unique genetic characteristics for specific grower needs and protected by U.S. patent(s) : PENDING.			





100088R14-61

CODE SEED-APPLIED TECHNOLOGY

- A Metalaxyl
- B Acceleron® B-300 SAT
- F Prothioconazole
- H1 Clothianidin  
at 0.25 mg ai/seed rate
- H2 Clothianidin  
at 0.50 mg ai/seed rate
- H3 Clothianidin  
at 1.25 mg ai/seed rate
- I Trifloxystrobin
- L Fluroxastrobin
- N Trifluralin
- V *Bacillus firmus* I-1582
- W Iproconazole



**WARNING: TREATED SEED. DO NOT USE TREATED SEED FOR FEED, FOOD OR OIL PURPOSES.**

This seed has been treated at the manufacturer's recommended rate with the products identified by code on the front of this seed tag, and decoded in the table in this section.

Clothianidin-treated seed: This seed has been treated at the rate indicated by code on the front of this seed tag, and decoded in the table in this section.

SEE INSIDE PANEL FOR TREATMENT SPECIFIC ADDITIONAL INFORMATION.

General Information: Handle treated seeds in ventilated areas. Avoid breathing or making contact with dust. If contaminated, remove clothing and launder before reuse.

Wash hands and face before eating, drinking, or smoking. Treated seed must be adequately covered with soil at planting. Treated seeds must be planted at a minimum depth of 0.5 inches. To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations.

**Personal Protective Equipment:** When opening this bag, or loading, pouring, or handling the treated seed or seed pieces, wear long-sleeved shirt, long pants, shoes, socks and chemical resistant gloves made out of barrier laminate, nitrile rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyethylene, polyvinyl chloride ≥14 mils, or viton ≥14 mils.

**Hazards to Humans and Animals:** Hazardous to humans if swallowed, inhaled or absorbed through skin. Treated seed is toxic to birds and mammals. Exposed seed is hazardous to birds and mammals that come in contact with treated seed. Cover or collect treated seeds that are spilled during loading and planting, and in areas such as row ends. After seeds and/or seed pieces have been planted in soil or other planting media, DO NOT enter or allow workers or other animals into areas with treated seeds during the restricted-entry interval of 12 hours, except that workers may enter the areas with treated seeds without restriction if there will be no worker contact with the soil/media surface.

**Storage and Disposal:** Store treated seed away from food and feed, and do not allow access by children, pets, or livestock. Do not reuse or refill empty seed bags. Surplus treated seed in empty treated seed containers should be stored or disposed according to local requirements. Dispose of all excess treated seed by burying away from bodies of water. Do not contaminate bodies of water when cleaning up excess treated seed or wash water of planting equipment. Dispose of them seed and seed packaging in accordance with local requirements. Left over treated seed may be double sown around the headland or burned away from water bodies in accordance with local requirements. Excess treated seed may be used for other agricultural production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agonomic products.

**FIRST AID:** Eyes: Hold eye open and rinse gently with water for 15 minutes. If wearing contact lenses, remove after first 5 minutes and continue rinsing. Skin or clothing: Remove contaminated clothing. Rinse skin immediately with plenty of water. Use soap if available. Swallowed: Call poison center or doctor immediately for treatment advice. Sip water if able to swallow. Do not induce vomiting unless told to do so by the poison center or doctor. Do not give anything by mouth to an unconscious person. Inhaled: Move to fresh air. If not breathing, call 911 or ambulance, and then give artificial respiration. Call poison center or doctor for further treatment advice. **FOR EMERGENCY MEDICAL TREATMENT INFORMATION, CALL COLLECT DAY OR NIGHT: (314) 694-4000.**

**PRECAUTIONARY INFORMATION CONTINUED ON INSIDE PANEL.**

Call 1-800-8 DENALB (1-800-833-5252) with any questions, feedback or additional seed needs.

**NOTICE:** See bag for limitation of warranties, liability and remedies.  
Monsanto Company - Attention Seed Box - 800 N. Lindbergh Blvd. - St. Louis, MO 63167

100088R14-61 NET WEIGHT PRINTED ON BAG ©2008-2017 MONSANTO COMPANY