Site KBC Wollocalule Miscallinus	VBC	
Planting	Miscanthus x giganteus was planted in May of 2008	Arlington Miscanthus x giganteus was planted in May of
Cover Crop		be re-planted in 2010 with the same rootstock
Tillage	None	None
Hanvest	None	None
ומועפטנ	Same as G5, G7, G9, and G10	No harvest in 2010
	Harvest so that 6 inches of stubble remain for wildlife cover. Harvest in fall after hard frost / senescence	
reruization	Nitrogen will be applied at 50 lb N acre ⁻¹ (56 kg N ha ⁻¹)	No N, P, or K in 2009
Pest Catrl	No P and K in 2010	
	miscanthus begins to grow.	PRE application of Roundup at 22 oz/A applied before miscanthus begins to grow.
	Preemergence application of Dual II Magnum 1.5 pts/A	Pre-emergence: Dual II Magnum 1.74 ha ⁻¹ (1.5 pts/ac)
	If needed: PRE or POST application of Drive or Paramount (quinclorac) 0.5 lb/A + atrazine 0.5 lb/A.	Post-emergence: 2,4-D ester 0.61 I ha ⁻¹ (8oz ac ⁻¹) as needed
Microplote	Postemergence application of 2,4-D ester as needed.	
and opioe	No-N strip. No N fertilizer applied to west 15 feet of plot (except east 15' for block 1 of KBS to avoid gas chambers).	No-N strip. No N fertilizer applied to west 15 feet of plot (not applicable in 2010)