

**GLYPHOSATE**  
Tolerance

**GLUFOSINATE**  
Tolerance

See important herbicide use restrictions on back of tag.



**PIONEER**<sup>®</sup>

BRAND · PRODUCTS

**P0306Q**

BLEND/MIXTURE

COMPARATIVE RM 103

**F14**



MATERIAL: P0306Q-NI07 BATCH: 4620825

KIND: Field Corn LOT NO: B3CON13003-0

TOL1 TESTED: NOV 2020

KERNELS/UNIT: 80,000

FERTILE CYTOPLASM

Hybrid: P0306CYFR (95% of Pure Seed)

Germ: 95% Origin: MI

Hybrid: IRC03-03LR (5% of Pure Seed)

Germ: 95% Origin: NE

Noxious Weed Per Pound: None Other Crop Seed: None  
Pure Seed: 99.5% Inert Matter: 0.5% Weed Seed: None

The seed is a Mixture under the state laws of AL, AK, AZ, CT, DE, FL, GA, ID, IL, KS, LA, KY, MA, MD, MS, NC, NE, NJ, NV, NY, OK, OR, TN, UT, VA and WI

See Bag/Tag/Invoice for important terms of purchase and limited license. Limited license is granted solely to produce a single crop of grain for feeding or processing with no rights for unauthorized propagation, seed multiplication and exportations is prohibited.



**ABOVE/BELOW**

HERCULEX<sup>®</sup> XTRA  
Insect Protection



**Fungicide, Insecticide, Nematicide  
Seed Treatment**



NOTICE: See important Product Use Information on back of tag.



**MAJOR COMPONENT:**

This seed is treated at the manufacturer's recommended rates with fludioxonil, metenoxam, thiabendazole, ethaboxam, ipconazole, and azoxystrobin fungicides, clothianidin (0.25 mg ai/seed) insecticide, chlorantraniliprole insecticide, and *Bacillus amyloliquefaciens* strains PTA-4838 and MBI-600. 02/20

**REFUGE COMPONENT:**

This seed is treated at the manufacturer's recommended rates with fludioxonil, metenoxam, thiabendazole, ethaboxam, ipconazole, and azoxystrobin fungicides, clothianidin (1.25 mg ai/seed) insecticide, *Bacillus firmus* I-1582, and *Bacillus amyloliquefaciens* strain MBI 600. 02/20

**CAUTION-TREATED SEED: DO NOT USE TREATED SEED FOR FEED, FOOD OR OIL PURPOSES. DO NOT ALLOW CHILDREN, PETS, OR LIVESTOCK TO HAVE ACCESS TO TREATED SEED. STORE AWAY FROM FOOD AND FEEDSTUFFS. DO NOT USE EMPTY SEED BAGS FOR ANY OTHER PURPOSE.** Excess treated seed may be used for ethanol production only if (1) byproducts are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol byproducts that are used for agronomic practice.

**Safety & Health:** Wear long-sleeved shirt, long pants, shoes and socks, and waterproof chemical resistant gloves made out of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils when handling treated seed. For more information contact provider at 1-800-932-5994. **Restricted Entry Interval:** After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction. **Environmental:** This compound is toxic to birds and mammals. Do not use in areas where rice/crawfish aquaculture are in place or in or near fish farms, shrimp, prawn or crab ponds or nursery operations. Treated seeds exposed on soil surface may be hazardous to birds, mammals, aquatic invertebrates, and wildlife. Store or collect treated seeds spilled during loading. Dispose of all excess treated seed. Left over treated seed may be double sown into ground water if used in areas where soils are permeable, particularly where the water table is shallow. **Additional Information:** Use of chlorantraniliprole as a seed treatment must be communicated to all personnel involved in seasonal insect control applications. Treated seed must be planted into the soil at a depth greater than 1 inch. Chlorantraniliprole may be applied to seed already treated with chlorantraniliprole. Following the planting of chlorantraniliprole-treated seed, do not apply chlorantraniliprole or other Group 28 products for approximately 60 days. Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide). Following the planting of chlorantraniliprole-treated seed, total foliar applications to a field corn crop may not exceed a total rate of 0.146 lb chlorantraniliprole per acre. Soil applications of chlorantraniliprole-containing products are not permitted to corn if chlorantraniliprole treated seeds are planted. Treated seed must be planted into the soil at a depth greater than 1 inch. Thiamethoxam: Do not make any soil or foliar applications of products containing thiamethoxam to corn crops grown from seed treated with thiamethoxam. Do not use at a rate that will result in more than 0.165 lb thiamethoxam per acre corn (74.8 grams ai/A) per season. Regardless of type of application (seed treatment, soil or foliar), do not apply more than 0.1 lb corn of active ingredient clothianidin per acre per year. Clothianidin is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert. **Grazing and Plantback:** Do not graze or feed soybean forage and hay to livestock. Corn forage may not be grazed until 30 days after planting. In the event of crop failure or harvest of a crop grown from seed treated with Maxim Quattro, crops may be replanted according to the following schedule: **Immediate:** Alfalfa, Cereal Grains: Barley, Corn, Oat, Rye, Triticale and Wheat, Cucurbit Vegetables Crop Group 9, Head and Stem Brassica Crop Group 9, SA, Onion, Bulb, Crop Subgroup 3-47A, Peas, Dried Shelled: Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea, Root Vegetables (except Sugarbeet) Crop Subgroup 18, Soybean, Spinach, Sweet Potato, For any other crops the minimum plantback interval is 30 days from the date seeds treated with Maxim Quattro were planted. **Thiamethoxam:** In event of a crop failure or harvest of thiamethoxam treated seed, the field may be replanted immediately to alfalfa, Brassica (cole) leafy vegetables, cereal grains (including barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice (dry-seeded), rye, sorghum, teosinte, triticale, wheat and wild rice), canola, cotton, cucurbit vegetables, dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables, mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, peanuts, rapeseed seed, and safflower seed), root vegetables, strawberry, sunflowers, tobacco, and tuberous and corn vegetables. For any other crop, the minimum plant back interval is 120 days from the date the thiamethoxam-treated seed was planted. A cover crop other than the crops listed that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however the crop may not be grazed or harvested for food. **Clothianidin:** Immediate plantback: cereal grains, field corn, popcorn, sweet corn, rapeseed, canola, cotton, cucurbit vegetables, dry bulb vegetables; 30 day plant back: grasses, non-grass animal feeds, dry beans, dry peas; 8 month plant back: sugar cane; 12-month plant back: any crops without an earlier plant back interval. **Chlorantraniliprole:** Crops on the Lumnia label and the following crops or crop groups may be planted immediately following harvest: Artichoke, Asparagus, Brassica (Cole) Leafy Vegetables (Crop Group 9); Corn (pop. seed, and sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Fruiting Vegetables (Crop Group 8); Grape; Grass Forage, Fodder and Hay (Crop Group 17); Herbs subgroup (Crop Group 19A); Leafy Vegetables (nonbrassica, Crop Group 4); Legume Vegetables (Crop Group 5); Foliage of Legume Vegetables (Crop Group 7); Spearmint; Peppermint; Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay Crop Group 18); Okra; Potato; Prickly Pear Cactus; Root and Tuber Vegetables (Crop Group 1); Strawberries; Sugarcane; Tobacco and Tops of Root and Tuber Vegetables (Crop Group 2). The following crops or crop groups may be planted 30 days following the last planting of chlorantraniliprole treated seed: garlic; greaththeaded garlic; dry bulb onion; leek; green onion; Welsh onion; shallot; Cereal Grains (Crop Group 15); soybean and peanut. All other crops cannot be planted until 12 months after the last planting of chlorantraniliprole treated seed. 02/20

**PRODUCT USE STATEMENT:** This seed is a blend of 5% refuge seed and 95% seed containing the Herculex I and Herculex RW Insect Protection genes that produce a *Bacillus thuringiensis* (*Bt*) Cry1F protein and the *Bt* Cry34Ab1 and Cry35Ab1 proteins; the Agrisure<sup>®</sup> RW trait that includes a gene that produces a *Bt* mCry3A protein, and the YIELDGARD<sup>®</sup> Corn Borer gene which produces a *Bt* Cry1Ab protein that provide protection or suppression against susceptible European corn borer, southwestern corn borer, black cutworm, fall armyworm, western bean cutworm, lesser corn stalk borer, southern corn stalk borer, stalk borer, sugarcane borer, and corn earworm; and also provide protection from larval injury caused by susceptible western corn rootworm, northern corn rootworm and Mexican corn rootworm. Product responses may vary by location, post population, environmental conditions, and agricultural practices. These proteins and the genetic material necessary for their production in corn are protected under one or more U.S. patents. **YOU MUST SIGN A TECHNOLOGY USE AGREEMENT, READ THE PRODUCT USE GUIDE PRIOR TO PLANTING AND FOLLOW INSECT RESISTANCE MANAGEMENT (IRM) REQUIREMENTS.**

**PATENT STATEMENT:** The Herculex I and Herculex RW Insect Resistance technologies incorporated into these seeds are protected under one or more U.S. patents. The purchase of these seeds includes a limited license to produce a single corn crop in the United States (or other applicable country). The use of seed from such a crop for the progeny thereof for propagation or seed multiplication or for production or development of a hybrid or different variety of seed is strictly prohibited. Agrisure RW technology incorporated into these seeds is commercialized under a license from Syngenta Crop Protection AG, under one or more U.S. patent numbers. This license does not extend to the use of seed from such crop or the progeny thereof for propagation or seed multiplication. Furthermore, the use of such seed or the progeny thereof for propagation, seed multiplication, production or development of a hybrid or different variety of seed, research, breeding or crossing, is strictly prohibited. Resale or transfer of the seed is strictly prohibited. **YIELDGARD** is a registered trademark used under license from Monsanto Co. **HERCULEX** is a registered trademark of Dow AgroSciences and Pioneer Hi-Bred. **Herculex** is a registered trademark of Dow AgroSciences LLC. **Agrisure** is a registered trademark of, and used under license from, a Syngenta Group company. **Agrisure** technology incorporated into these seeds is commercialized under a license from Syngenta Crop Protection AG. US Crome 7-19

**PRODUCT USE STATEMENT:** This hybrid contains the Roundup Ready<sup>®</sup> Corn 2 gene developed by Monsanto Company. The Roundup Ready<sup>®</sup> gene provides resistance to glyphosate, the active ingredient in labeled Roundup<sup>®</sup> branded herbicides. **WARNING:** The Roundup Ready gene will safeguard this hybrid **ONLY** against applications of glyphosate. The active ingredient in labeled Roundup branded herbicides, when applied at labeled rates, The Roundup Ready gene **WILL NOT** safeguard this hybrid against applications of other herbicides which require a different herbicide resistance gene. Always read and follow herbicide label directions prior to use. **ACCIDENTAL APPLICATIONS OF INCOMPATIBLE HERBICIDES TO THIS HYBRID COULD RESULT IN TOTAL CROP LOSS.**

**YOU MUST SIGN A TECHNOLOGY USE AGREEMENT, READ THE PRODUCT USE GUIDE PRIOR TO PLANTING THIS SEED IS ACQUIRED UNDER AN AGREEMENT THAT INCLUDES THE FOLLOWING TERMS:** The following United States patents: 6,826,400; 7,582,434; 8,273,959 and 8,722,969. The purchase of these seeds includes a limited license to produce a single corn crop in the United States (or other applicable country). This license does not extend to the use of seed of such crop or the progeny thereof for propagation or seed multiplication or breeding and the foregoing is strictly prohibited. **Roundup** and Roundup Ready are registered trademarks of Monsanto Technology LLC. RPS2 Corn Rev. 7-19

**PRODUCT USE STATEMENT:** This hybrid contains the LibertyLink<sup>®</sup> gene. These seeds and the plants grown from these seeds produce a PAT (phosphinothricin acetyltransferase) protein that provides resistance to the Liberty<sup>®</sup> Herbicide (glufosinate). **WARNING:** The LibertyLink gene will safeguard this hybrid **ONLY** against applications of Liberty. The LibertyLink gene **WILL NOT** safeguard this hybrid against applications of other herbicides which require a different herbicide resistance gene. Always read and follow herbicide label directions prior to use. **ACCIDENTAL APPLICATIONS OF INCOMPATIBLE HERBICIDES TO THIS HYBRID COULD RESULT IN TOTAL CROP LOSS.** **YOU MUST SIGN A TECHNOLOGY USE AGREEMENT, READ THE PRODUCT USE GUIDE PRIOR TO PLANTING.** The herbicide resistance technology contained in this seed is protected at least under one or more of the following U.S. patents: RF4-0,967. The conditional purchase of these seeds conveys no license under said patents to use these seeds, except to produce a single commercial crop (one time use only) in the United States, and harvesting the grain solely for purposes of food or feed applications or for industrial processing. This license does not extend to the use of seed of such crop or the progeny thereof for propagation or seed multiplication or breeding and the foregoing is strictly prohibited. **LibertyLink**, Liberty<sup>®</sup> and the Water Droplet Design are registered trademarks of BASF. RPS2 Corn Rev. 7-19

Pioneer<sup>®</sup> brand products are provided subject to the terms and conditions of purchase. <sup>®</sup>, <sup>™</sup> Trademarks and service marks of DuPont, Pioneer or their respective owners. © 2017 PHIL. PIONEER HI-BRED INTERNATIONAL, INC., PO BOX 256, Johnston, IA 50131-0256




**IN CORN-GROWING AREAS, THE FOLLOWING REFUGE REQUIREMENTS MUST BE FOLLOWED:**

- NO ADDITIONAL REFUGE IS NEEDED IN CORN-GROWING AREAS.
- Read and follow Product Use Guide for foliar insecticide application requirements if economic thresholds for certain target pests are reached.


**IN COTTON-GROWING AREAS, THE FOLLOWING REFUGE REQUIREMENTS MUST BE FOLLOWED:**

- Plant at least 20% of the corn acres to a corn borer refuge in COTTON-GROWING AREAS.
- The refuge should be sown on the same day, or with the shortest window possible between planting dates.
- External refuges may be planted as an in-field or adjacent (e.g., across the road) refuge or as a separate block within 1/2 mile of the integrated refuge field.
- In-field refuge options include: blocks, perimeter strips (i.e., along the edges or headlands), or in-field strips.
- When planting the refuge in strips across the field, refuges must be at least four (4) rows wide.
- Refuge hybrids must not express a Bt protein active against corn borer.
- Read and follow Product Use Guide for foliar insecticide application requirements if economic thresholds for certain target pests are reached.
- For full details, consult the Product Use Guide.

**In-field Refuge**



**Adjacent Field Refuge**



**Key** | ■ Bt Product   ■ ≥ 20% Corn Borer Refuge



**Cotton-Growing Areas are all the counties in:**  
AR, LA, MS, AL, GA, FL, SC, NC and in:

MO counties of Dunklin, New Madrid, Pemiscot, Scott and Stoddard

OK counties of Beckham, Caddo, Comanche, Custer, Greer, Harmon, Jackson, Kay, Kiowa, Tillman and Washita

TN counties of Carroll, Chester, Crockett, Dyer, Fayette, Franklin, Gibson, Hardeman, Hardin, Haywood, Lake, Lauderdale, Lincoln, Madison, Obion, Rutherford, Shelby and Tipton

TX counties except Carson, Dallam, Hansford, Hartley, Hutchinson, Lipscomb, Moore, Ochiltree, Roberts and Sherman

VA counties of Dinwiddie, Franklin City, Greensville, Isle of Wight, Northampton, Southampton, Suffolk City, Surrey and Sussex

For additional refuge planning tools, you may also log onto [www.irmcalculator.com](http://www.irmcalculator.com).

**NO ADDITIONAL REFUGE NEEDED**

Corn-growing Areas

**20% REFUGE**  
Cotton-growing Areas

**CORN BORER REFUGE OPTIONS**

38806  
18D-1211 Integrated Refuge