

# EQUITY VIP

This seed has been treated with thiamethoxam insecticide and fludioxonil fungicide, thiazoxazole and seedborne fungicides.

Do not use for feed, food, or oil purposes

#### Ground Water Advisory:

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow. Methoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### Pollinator Precautions:

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.

In the event of a crop failure or harvest of a crop grown from THX/MXM/FLD/7BZ/SDX FS treated seed, the field may be replanted according to the following schedule:

Immediate Plantback	Minimum 30-Day Plantback Interval	Minimum 120-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat Peas, Dried Shelled Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea Soybean	Alfalfa Brassica (Cauli) Leafy Vegetables Cereal Grains (buckwheat, corn, pearl millet, proso, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice Cotton Cucurbit Vegetables Fruiting Vegetables Globe Artichoke Leafy Vegetables Legume Vegetables (succulent or dried) Mint, Peppermint and Spearmint Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower Peanut Root Vegetables Strawberry Sunflower Tobacco	All Other Crops

For any other crop, the minimum plant back interval is 120 days from the date the THX/MXM/FLD/7BZ/SDX FS treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval, however, the crop may not be grazed or harvested for food or feed.

Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.3 grams a/A) per acre, 0.004 lb fludioxonil (2.0 grams a/A) per acre and 0.013 lb metoxam (5.7 grams a/A) per acre per calendar year as a seed treatment application.

This seed has been treated with 0.0756 mg of thiamethoxam, 0.0113 mg metoxam, and 0.0039 mg fludioxonil per seed. Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with THX/MXM/FLD/7BZ/SDX.

7339 D0116

  
**Loveland**  
 PRODUCTS  
 Loveland Products, Inc. • P.O. Box 1286 • Greeley, CO 80632

# DYNA-START™ 3P™

**Dyna-Start 3P+ Single Strain Soybean  
Inoculant + Dyna-Start 3P+ Biological +  
Consensus**

This seed has been treated with ActivAC™-tive Inge Ingredients Indole-3-bredients Indole-3-butyryltryric acid, Chitosan, and Salicylic Acid atacid, Chitosan, and Salicylic Acid at the rate specified bate specified by they the manufacturer.

This seed has been treated with Bradyrhizobium japonicum at the rate specified by the manufacturer.

This seed has been treated with Bacillus Amylolyticus at the rate specified by the manufacturer.

Not Food use.

Do not use treated seed for for food, oil or for feed purposes.

Store treated seed away from food and hand feedstuffs.

Do not allow children, pets or children, pets or livestock to have access to treated access to treated seeds.

Dispose of seed packaging or containers in accordance containers in accordance with with local requirements.

  
**Loveland**  
 PRODUCTS

FORMULATED FOR: Loveland Products, Inc., P.O. Box 1286, Greeley, Colorado 80632-1286  
 Copyright © 2023 Loveland Products, Inc. All Rights Reserved.  
 CONSENSUS™ and DYNA-START™ are registered trademarks of Loveland Products, Inc.